

# Huaier aqueous extract suppresses human breast cancer cell proliferation through inhibition of estrogen receptor $\alpha$ signaling

XIAOLONG WANG, NING ZHANG, QIANG HUO, MINGJUAN SUN, SHANGGE LV and QIFENG YANG

Department of Breast Surgery, Qilu Hospital, Shandong University, Jinan, Shandong 250012, P.R. China

DOI: 10.3892/ijo\_XXXXXXXX

**Abstract.** Estrogen receptor  $\alpha$  (ER $\alpha$ ) has been reported to play a critical role in promoting the growth of breast tumor cells. In the present study, we explored the effect of Huaier extract on estrogen receptor  $\alpha$  signaling in breast cancer cell lines. Our data demonstrated that Huaier extract effectively inhibited the proliferation of the MCF-7, T47D and ZR-75-1 human breast cancer cell lines. For the mechanism analysis, we demonstrated that Huaier extract significantly reduced the mRNA and protein levels of ER $\alpha$  in all three ER $\alpha$ -positive cell lines. The downregulation of ER $\alpha$  protein levels was correlated with activation of the proteasomes. We demonstrated that Huaier extract markedly decreased the expression of both ER $\alpha$  and its downstream genes, inhibited the estrogen-stimulated proliferation and reversed the estrogen-induced activation of the nuclear factor  $\kappa$ B (NF $\kappa$ B) pathway. Our study provides evidence that Huaier extract is a novel estrogen receptor modulator and is a promising drug for the prevention and treatment of ER $\alpha$ -positive human breast cancers.

## Introduction

Estrogen receptors (ER $\alpha$  and ER $\beta$ ) are members of the superfamily of nuclear steroid hormone receptors (1). After the binding of hormone to these receptors, the hormone-receptor complexes directly bind to specific sequences on the DNA, named estrogen response elements (EREs), or interact with other transcription factors (2-4). ER $\alpha$  and ER $\beta$  are expressed both in normal and cancerous breast epithelium (5). It has been reported that a high ER $\alpha$ :ER $\beta$  ratio was responsible for enhanced cell proliferation (6). Consistent with the critical role of ER $\alpha$  in mediating the proliferative effect of estrogen, mice that selectively lacked ER $\alpha$  showed a severely under-developed mammary epithelium with only rudimentary ductal structure (7,8). In 2003, Shaaban *et al* demonstrated that the malignant progression of breast lesion was correlated with elevated ER $\alpha$  expression (9). Therefore,

selective disruption of ER $\alpha$  signaling in breast cancer cells may be a potent therapy to prevent the development of breast cancers.

Current selective estrogen receptor modulators (SERMs) exhibit their effects through three different ways (10): i) interfering with estrogen receptor  $\alpha$  (ER $\alpha$ ) binding, such as tamoxifen and faslodex; ii) inhibiting endogenous estrogen production, such as oophorectomy, anastrozole and letrozole; and iii) down-regulating ER $\alpha$  protein level, such as fulvestrant. SERMs may exert estrogen agonist action in some target tissues while acting as estrogen agonist action in others (11). Therapeutical SERMs are considered to be able to inhibit the proliferation of breast cancer without the influence on bone density, cholesterol level or uterine endometrium (12,13).

Natural products have long been considered as a rich source of novel bioactive chemicals providing potential effective new drugs. Epidemiological and physiological studies have shown that phytochemicals from vegetables and fruits possess potent anticancer activities with minimal to no side-effect (14). Huaier extract is a kind of officinal fungi and has been used in China for nearly 1,600 years. In the past few years, its anticancer activities are attracting increasing interest. Our previous studies showed that the antitumor effect of Huaier extract on breast cancer may be associated with induction of apoptosis and inhibition of angiogenesis (15,16). Similar effects were seen in human liver cancer (17-19). In view of these findings, it was of great interest to determine whether Huaier extract could affect the ER signaling pathway in human breast cancer cells.

In this study, we used three ER $\alpha$ -positive breast cancer cell lines (MCF-7, T47D and ZR-75-1). Our findings demonstrated that Huaier extract markedly decreased expression of both ER $\alpha$  and its downstream genes, inhibited the estrogen-stimulated proliferation and reversed the estrogen-induced activation of the NF- $\kappa$ B pathway. These effects provided a further rationale for the use of Huaier extract in the prevention and treatment of ER $\alpha$ -dependent human breast cancers.

## Materials and methods

**Cell cultures and reagents.** Human breast cancer cell lines, MCF-7, T47D and ZR-75-1, were purchased from ATCC (Manassas, VA, USA). MCF-7 cells were routinely maintained in DMEM medium (Gibco-BRL, Rockville, IN, USA) supplemented with 10% fetal bovine serum (Haoyang Biological Manufacture Co. Ltd., Tianjin, China), 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin. T47D cells were cultured in RPMI-1640 medium

*Correspondence to:* Dr Qifeng Yang, Department of Breast Surgery, Qilu Hospital, Shandong University, Wenhua Xi Road no. 107, Jinan, Shandong 250012, P.R. China  
E-mail: qifengy@gmail.com

**Key words:** Huaier, estrogen receptor, nuclear factor  $\kappa$ B, breast cancer




Figure 1. Huaier extract inhibits the proliferation of ER $\alpha$ -positive human breast cancer cells. (A) MCF-7, (B) T47D and (C) ZR-75-1 were exposed to various concentrations of Huaier extract for 24, 48 or 72 h. Data shown are averages of triplicate experiments, and error bars represent SD. \*P<0.05 vs. control group.

(Gibco-BRL) with 10% fetal bovine serum and 10  $\mu$ g/ml bovine insulin (Sigma-Aldrich, St. Louis, MO, USA). ZR-75-1 cells were cultured in the above medium in the absence of bovine insulin. For estrogen-free experiments, medium was changed to phenol red-free DMEM or RPMI-1640 with 0.5% dextran-coated charcoal (DCC) stripped FBS (Haoyang Biological Manufacture) for 72 h. The medium was then changed to the indicated serum and phenol red-free medium for 24 h before various treatments as described in the figure legends (20,21). Electuary ointment of Huaier extract was kindly provided by Gaitianli Medicine Co. Ltd. (Jiangsu, China). The stock solutions were generated by dissolving electuary ointment in indicated medium (15,16). MG 132 was purchased from Sigma-Aldrich.  $\beta$ -estradiol (E2) was obtained from Wako and was dissolved in ethanol with a final concentration of ethanol in medium of less than 0.1%. Anti-ER $\alpha$  antibody was from Dako. All other antibodies were purchased from Cell Signaling Technology.

*3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay.* Cells were plated into 96-well plates and grown overnight at 37°C in 5% CO<sub>2</sub> for 24 h. After treatment with the appropriate concentration of Huaier extract for indicated time, 20  $\mu$ l of MTT solution (5 mg/ml) was added to each well and incubated for 4–6 h at 37°C. The formazan crystal was dissolved in DMSO for 10 min. Optical density (OD) was measured by the Microplate Reader (Bio-Rad, Hercules, CA, USA).

For E2-stimulated proliferation, cells were seeded in 96-well plates in phenol red-free medium in the presence of 10% charcoal-stripped fetal bovine serum. The next day, the medium was changed to fresh serum and phenol red-free

medium for another 24 h. Then 10 nM E2 in the presence or absence of different concentrations of Huaier extract was added for 48 h.

*Western blot analysis.* For western blot analysis, treated cells were washed with ice-cold PBS and lysed in a modified RIPA buffer (1X PBS, 1% NP-40, 0.1% sodium dodecyl sulfate, 5 mM EDTA, 0.5% sodium deoxycholate and 1 mM sodium orthovanadate) with protease inhibitors for 20 min at 4°C. After centrifugation at 12,000 rpm for 15 min at 4°C, the supernatant was collected. Subsequently, 50  $\mu$ g of total cellular protein from each sample were separated by 10% SDS-PAGE and electrotransferred onto a polyvinylidene fluoride (PVDF) membranes by a semi-dry blotting apparatus (Bio-Rad). Then the membranes were immunoblotted with primary antibodies followed by horseradish peroxidase-conjugated secondary antibodies (1:7,000). Labeled protein spots were visualized by ECL (PerkinElmer) according to the manufacturer's instruction.  $\beta$ -actin was used as the loading control.

*Quantitative real-time reverse transcriptase (RT)-PCR.* Total RNA was isolated from cells using the TRIzol reagent (Invitrogen, Carlsbad, CA, USA) and methods described previously (22). Briefly, total RNA from treated cells were isolated with TRIzol reagent according to the manufacturer's protocol. Total RNA was used to synthesize cDNA using PrimerScript RT Reagent kit (Takara). The cDNA reaction product was amplified with following primers: ER $\alpha$  forward, 5'-CGGCATTCTACAGGCCAAATT-3' and ER $\alpha$  reverse, 5'-AGCGAGTCTCCTTGGCAGATT-3'; PR forward, 5'-AGCC




Figure 2. Huaier extract significantly decreases the mRNA levels of ER $\alpha$  in all three ER $\alpha$ -positive human breast cancer cell lines. (A) MCF-7, (B) T47D and (C) ZR-75-1 cells were treated with various concentrations of Huaier extract for indicated time. The mRNA levels were detected through real-time PCR. GAPDH was used as internal control for normalization. Data are representative of three independent experiments. \*P<0.05.

CACAATACAGCTTCGAG-3' and PR reverse, 5'-TTTCGAC CTCCAAGGACCAT-3'; pS2 forward, 5'-TGACTCGGGGTC GCCTTTGGAG-3' and pS2 reverse, 5'-GTGAGCCGAGGC ACAGCTGCAG-3'; cathepsin D forward, 5'-TGAGGCCATT GTGGACAAGGCAC-3' and cathepsin D reverse, 5'-GTCACG GTCAAACACAGTGTAGTAG-3'; GAPDH forward, 5'-AGAA GGCTGGGGCTCATTTG-3' and GAPDH reverse, 5'-AGGG GCCATCCACAGTCTTC-3'. GAPDH was used for equal RNA loading. The experiments were repeated for at least three times.

**Statistical analysis.** Data were expressed as means  $\pm$  standard deviations (SD) for three replicate experiments and considered significant at P<0.05. The statistical analysis was carried out by using SPSS edition 19.0.

## Results

**Huaier extract effectively inhibited the proliferation of ER $\alpha$ -positive breast cancer cell lines.** We first examined the effect of Huaier extract on the growth of three ER $\alpha$ -positive human breast cancer cell lines, MCF-7, T47D and ZR-75-1. As

shown in Fig. 1, Huaier extract significantly inhibited the proliferation of these three cell lines in a dose- and time-dependent manner (P<0.05). The results from MCF-7 were similar to our previous reports (15). In addition, at 24 h, the respective IC<sub>50</sub> values for MCF-7 (Fig. 1A), T47D (Fig. 1B) and ZR-75-1 (Fig. 1C) were 7.27 $\pm$ 0.86, 5.12 $\pm$ 0.71 and 1.30 $\pm$ 0.11 mg/ml. Thus the ZR-75-1 cell line was more sensitive to the growth inhibition caused by Huaier extract than the other two.

**Huaier extract effectively decreased the mRNA and protein levels of ER $\alpha$  in three ER $\alpha$ -positive human breast cancer cell lines.** ER $\alpha$  is generally associated with cell cycle progression and has been reported to promote both anchorage-dependent and anchorage-independent growth of breast cancer cells (23-25). To determine the effect of Huaier extract on the level of ER $\alpha$ , cell lines were exposed to various concentrations of Huaier extract for 24, 48 or 72 h. We then examined the effect of Huaier extract on ER $\alpha$  mRNA levels. As shown in Fig. 2A, Huaier extract could decrease the mRNA level of ER $\alpha$  dose- and time-dependently in MCF-7 cell line. After various times of 4 mg/ml Huaier treatment, the transcriptional levels were significantly inhibited




Figure 3. Huaier extract significantly reduces the protein levels of ER $\alpha$  through activated proteasome pathway. ER $\alpha$  levels were examined by western blot assay. Actin was used as a loading control. (A-C) After treatment with Huaier extract, ER $\alpha$  levels in MCF-7, T47D and ZR-75-1 were reduced in a dose- and time-dependent manner. (D and E) MCF-7 and T47D cells were treated with vehicle, 4 mg/ml Huaier extract or 10  $\mu$ m MG 132 in combination with Huaier extract for 24 h. MG 132 was added 1 h before the Huaier extract treatment. The bar graphs represent the mean relative densities  $\pm$  SD of three independent experiments. \*P<0.05 with respect to the basal conditions. <sup>#</sup>P<0.05 with respect to Huaier treatment alone.

by 32.9 $\pm$ 2.7, 55.4 $\pm$ 4.1 and 72.7 $\pm$ 6.3% (P<0.05), respectively. Reduction of ER $\alpha$  mRNA levels was also observed in T47D and ZR-75-1 cells (Fig. 2B and C).

Data from Fig. 3 demonstrated that Huaier extract markedly downregulated the ER $\alpha$  protein levels in a time- and dose-dependent manner (P<0.05). After 24 h incubation of 4 mg/ml Huaier extract, the levels of ER $\alpha$  were reduced by 67.2 $\pm$ 9.8, 86.1 $\pm$ 7.0 and 85.3 $\pm$ 4.0% in MCF-7 (Fig. 3A), T47D (Fig. 3B) and ZR-75-1 (Fig. 3C) cell lines, respectively. Reduction of ER $\alpha$  protein levels was observed in all three ER $\alpha$ -positive breast cancer cell lines, suggesting that the impact of Huaier extract on this protein is common to ER $\alpha$ -positive breast cancer cell lines.

*Huaier-induced downregulation of ER $\alpha$  protein was associated with activated proteasome.* Previous studies have revealed that regulation of ER $\alpha$  protein is associated with protein degradation

via proteasomes (26). To explore the involvement of proteasome in the Huaier-induced ER $\alpha$  degradation, the proteasome inhibitor, MG132, was used. Treatment of MCF-7 cells with Huaier extract led to a potent and persistent decrease in ER $\alpha$  protein levels. Addition of MG132 before Huaier treatment markedly rescued the downregulation of ER $\alpha$  (P<0.05, Fig. 3D). In addition, MG132 was also able to abolish the inhibitory effect of Huaier on ER $\alpha$  protein in T47D cell line (Fig. 3E). These findings demonstrated that inhibition of both gene transcription and proteasome-mediated degradation was required to reduce endogenous levels of ER $\alpha$  in response to Huaier extract treatment.

*Huaier extract significantly downregulates the transcription of estrogen responsive genes.* It has been reported that some activators of ER $\alpha$ , such as 17- $\beta$ -estradiol, could also decrease ER $\alpha$  level (27), suggesting that a reduced level of ER $\alpha$  was not




Figure 4. Huaier extract reduces the levels of target genes regulated by estrogen receptor. (A) MCF-7 and (B) ZR-75-1 cells were treated with various concentrations of Huaier extract for 10, 19 or 24 h. mRNA levels of cathepsin D, pS2 and PR were determined by real-time PCR as described in Materials and methods. Data are representative of three independent experiments. \* $P < 0.05$ .

closely associated with a reduction of ER $\alpha$  activity. Therefore, we next determined the transcriptional activity of ER $\alpha$  by detecting its downstream target genes after Huaier extract treatment (28,29).

The expressions of cathepsin D, pS2 and progesterone receptor (PR) were reported to be regulated by estrogen (30). To further determine the transcriptional activity of ER $\alpha$ , we examined their mRNA levels after Huaier extract treatment within 24 h. The levels of transcription of each gene were normalized to the level of GAPDH. As shown in Fig. 4A, MCF-7 cells were cultured in the absence or presence of Huaier extract (2-8 mg/ml) for 10, 19 or 24 h. Significant reduction of mRNA could be observed dose- and time-dependently ( $P < 0.05$ ). Similar results were obtained in ZR-75-1 cells (Fig. 4B). However, as PR expression in T47D cells was demonstrated to be independent of estrogen (31), we did not detect the mRNA level of PR in T47D cells. In addition, Huaier extract failed to cause significant reduction on gene expression of pS2 and cathepsin D in T47D cells (data not shown).

**Huaier extract inhibits E2-induced proliferation and NF- $\kappa$ B activation.** The proliferation of ER $\alpha$ -positive human breast cancer cell lines is strongly stimulated by estrogens (25). To specifically determine the inhibitory effect of Huaier extract on E2-induced effect, MTT and western blot assays were used. As shown in Fig. 5A, Huaier extract significantly inhibited

E2-induced cell proliferation in a dose-dependent manner. The survival of MCF-7 cells was significantly reduced by  $49.1 \pm 4.6\%$  after 48-h treatment with 1 mg/ml Huaier extract as compared with E2 treatment alone. The inhibition was observed in both MCF-7 and T47D cell lines (Fig. 5A and B).

It has been reported that activation of NF- $\kappa$ B contributed to the estrogen-induced proliferation in breast cancer cells (32). Thus, we detected the effect of Huaier extract on NF- $\kappa$ B pathway. The activation of NF- $\kappa$ B was assessed at 1 h in MCF-7 and T47D cells treated with control, estrogen alone or in combination with Huaier extract (2-4 mg/ml). As shown in Fig. 5C and D, 10 nM estrogen significantly phosphorylated P65 without influencing the total protein level of P65. After addition of Huaier extract, the phosphorylated levels of P65 were reduced almost to the basal level without estrogen. These results demonstrated that Huaier extract completely abolished the effect of estrogen on the activation of NF- $\kappa$ B and thus suppressed the proliferation of breast cancer cells induced by estrogen.

## Discussion

ER $\alpha$  has become an important target in the treatment of hormone-responsive breast cancers. It has been reported that treatment with tamoxifen has enhanced patient survival (33). Unfortunately, most patients initially responding to anti-estrogen therapies, such as tamoxifen, will eventually




Figure 5. Huaier extract inhibits the cell growth and activation of NF- $\kappa$ B pathway stimulated by estrogen. (A and C) MCF-7 and (B and D) T47D cells were maintained in the serum- and phenol red-free medium in the absence or presence of 10  $\mu$ M E2 alone or in combination with different concentrations of Huaier extract. (A and B) Cell viability was determined using MTT assay after incubation of the above medium for 48 h. (C and D) Western blot assay was performed after presence of the above medium for 1 h. The results were obtained from three independent experiments and are shown as the mean  $\pm$  SD. \* $P$ <0.05 with respect to the basal conditions. # $P$ <0.05 with respect to E2 treatment alone.

become resistant (34), along with a possible association with endometrial carcinomas (35,36). Although the mechanisms of endocrine resistance are not fully identified, cross-talk between ER $\alpha$  and growth factor signaling pathways may be involved (37,38). Therefore, downregulating the levels of ER $\alpha$  may be a potent therapeutic therapy for both primary estrogen-dependent breast cancer and hormone-refractory breast cancer.

Fulvestrant (ICI 182780) is a selective estrogen receptor downregulator (SERD) with no agonist effects and has been demonstrated to decrease the level of ER $\alpha$  protein to block both ligand-dependent and -independent receptor activation (39). However, recent studies have reported that fulvestrant could cause gastro-intestinal disturbance and hot flashes (40). Thus, alternative interventions such as compounds from natural products are needed to replace or to supplement current therapies.

Recently, Huaier extract was shown to exhibit antitumor and anti-angiogenesis activities both *in vitro* and *in vivo* (15,16). Studies in our laboratory have demonstrated that Huaier extract inhibited breast cancer growth via a direct pro-apoptotic effect

on tumor cells, as well as through an indirect effect on endothelial cells (15,16). However, little is known about the effect of Huaier extract on the estrogen receptor signaling pathway in breast cancers.

In the present study, we explored the mechanisms of action of Huaier extract on breast cancer cell growth by focusing on ER $\alpha$ . Following the treatment with various concentrations of Huaier extract, the mRNA and protein levels of ER $\alpha$  were significantly decreased in MCF-7, ZR-75-1 and T47D cell lines in a time- and dose-dependent manner (Figs. 2 and 3), suggesting that this effect was general in ER $\alpha$ -positive breast cancer cells. In search for detail mechanisms causing reduced ER $\alpha$  level, we observed that MG 132, a proteasome inhibitor, could effectively suppress Huaier-induced ER $\alpha$  degradation (Fig. 3D and E). Thus, our data demonstrated that Huaier extract reduced the mRNA levels of ER $\alpha$  and decreased its protein through promotion of the proteasome pathway. However, fulvestrant reduce ER $\alpha$  levels only through increasing protein turnover without affecting its mRNA levels (39). In this view, Huaier extract may be useful in the treatment of breast cancer that are resistant to the pure antiestrogen.

1 After binding to ER $\alpha$ , the estrogen-ER $\alpha$  complex will trans-  
 2 locate to the target DNA binding site, the estrogen responsive  
 3 element, in the promoter region of the target gene for gene  
 4 transcription activation (41). Therefore, we next examined the  
 5 expression level of downstream genes regulated by estrogen,  
 6 such as cathepsin D, pS2 and PR (30). After treatment by  
 7 Huaier extract, the mRNA level of these genes were signifi-  
 8 cantly reduced in a time- and dose-dependent manner (Fig. 4),  
 9 suggesting that the Huaier extract suppressed the transcription  
 10 activity of ER $\alpha$  along with its reduced levels.

11 To further examine the effect of Huaier extract on the  
 12 17 $\beta$ -estradiol-stimulated cell growth and its potential mecha-  
 13 nisms, we analyzed several pathways regulated by estrogen and  
 14 discovered NF- $\kappa$ B pathway. Although some studies showed  
 15 that estrogen inhibited the tumor necrosis factor- $\alpha$ -induced  
 16 activation of NF- $\kappa$ B (42), Rubio *et al* demonstrated no antago-  
 17 nism between ER and NF- $\kappa$ B in T47D and HC11 cells (32). In  
 18 addition, in their study, activation of NF- $\kappa$ B was needed for  
 19 estrogen-induced proliferation and expression of cyclin D1. In  
 20 our study, data from Fig. 5 shows that 17 $\beta$ -estradiol significantly  
 21 enhanced the phosphorylation of P65, the NF- $\kappa$ B component, in  
 22 a phenol red and serum-free condition. After addition of Huaier  
 23 extract, the level of P-P65 was reduced to the basal level without  
 24 estrogen. Therefore, we suggest that Huaier extract inhibited the  
 25 E2-induced proliferation of breast cancer through activation of  
 26 the NF- $\kappa$ B pathway.

27 In summary, we found that Huaier extract potently inhib-  
 28 ited the proliferation of ER $\alpha$ -positive breast cancer cells, and  
 29 identified ER $\alpha$  as a possible target for Huaier extract treatment.  
 30 Therefore, it is of great importance to develop this natural  
 31 product to treat or prevent ER $\alpha$ -positive breast cancers.

### 32 Acknowledgements

33 This study was supported by National Natural Science  
 34 Foundation of China (nos. 81072150, 81172529 and 81272903)  
 35 and Shandong Science and Technology Development Plan  
 36 (no. 2012GZC22115). We thank Shi Yan and Cunzhong Yuan  
 37 for technical support with experiments. We also thank Weiting  
 38 Gu and Min Gao for critical discussions and Dong Lun for  
 39 substantial help.  
 40  
 41  
 42

### 43 References

- 44 1. Hedden A, Müller V and Jensen EV: A new interpretation of  
 45 antiestrogen action. *Ann NY Acad Sci* 761: 109-120, 1995.
- 46 2. Klinge CM: Estrogen receptor interaction with estrogen  
 47 response elements. *Nucleic Acids Res* 29: 2905-2919, 2001.
- 48 3. Kushner PJ, Agard DA, Greene GL, *et al*: Estrogen receptor  
 49 pathways to AP-1. *J Steroid Biochem Mol Biol* 74: 311-317,  
 50 2000.
- 51 4. Safe S: Transcriptional activation of genes by 17 $\beta$ -estradiol  
 52 through estrogen receptor-Sp1 interactions. *Vitam Horm* 62:  
 53 231-252, 2001.
- 54 5. Dotzlaq H, Leygue E, Watson PH and Murphy LC: Expression  
 55 of estrogen receptor- $\beta$  in human breast tumors. *J Clin Endocrinol*  
 56 *Metab* 82: 2371-2374, 1997.
- 57 6. Ali S and Coombes RC: Estrogen receptor alpha in human  
 58 breast cancer: occurrence and significance. *J Mammary Gland*  
 59 *Biol Neoplasia* 5: 271-281, 2000.
- 60 7. Korach KS, Couse JF, Curtis SW, *et al*: Estrogen receptor gene  
 disruption: molecular characterization and experimental and  
 clinical phenotypes. *Recent Progr Horm Res* 51: 159, 1996.
8. Bocchinfuso WP and Korach KS: Mammary gland develop-  
 ment and tumorigenesis in estrogen receptor knockout mice.  
*J Mammary Gland Biol Neoplasia* 2: 323-334, 1997.
9. Shaaban AM, O'Neill PA, Davies MPA, *et al*: Declining  
 estrogen receptor-[beta] expression defines malignant progres-  
 sion of human breast neoplasia. *American J Surg Pathol* 27:  
 1502-1512, 2003.
10. Gralow JR: Optimizing the treatment of metastatic breast  
 cancer. *Breast Cancer Res Treat* 89: 9-15, 2005.
11. Shang Y and Brown M: Molecular determinants for the tissue  
 specificity of SERMs. *Science* 295: 2465-2468, 2002.
12. Osborne CK, Zhao HH and Fuqua SAW: Selective estrogen  
 receptor modulators: structure, function, and clinical use. *J Clin*  
*Oncol* 18: 3172-3186, 2000.
13. McCarthy TL, Clough ME, Gundberg CM and Centrella M:  
 Expression of an estrogen receptor agonist in differentiating  
 osteoblast cultures. *Proc Natl Acad Sci USA* 105: 7022-7027,  
 2008.
14. Cassady JM, Baird WM and Chang CJ: Natural products as a  
 source of potential cancer chemotherapeutic and chemopreven-  
 tive agents. *J Nat Prod* 53: 23-41, 1990.
15. Zhang N, Kong X, Yan S, Yuan C and Yang Q: Huaier aqueous  
 extract inhibits proliferation of breast cancer cells by inducing  
 apoptosis. *Cancer Sci* 101: 2375-2383, 2010.
16. Wang X, Zhang N, Huo Q and Yang Q: Anti-angiogenic and  
 antitumor activities of Huaier aqueous extract. *Oncol Rep* 28:  
 1167-1175, 2012.
17. Ren J, Zheng C, Feng G, *et al*: Inhibitory effect of extract of  
 fungi of Huaier on hepatocellular carcinoma cells. *J Huazhong*  
*Univ Sci Technolog Med Sci* 29: 198-201, 2009.
18. Xu X, Wei Q, Wang K, *et al*: Anticancer effects of Huaier are  
 associated with down-regulation of P53. *Asian Pac J Cancer*  
*Prev* 12: 2251-2254, 2011.
19. Sun Y, Sun T, Wang F, *et al*: A polysaccharide from the fungi  
 of Huaier exhibits anti-tumor potential and immunomodulatory  
 effects. *Carbohydr Polym* 92: 577-582, 2013.
20. Lippman M, Bolan G and Huff K: The effects of estrogens and  
 antiestrogens on hormone-responsive human breast cancer in  
 long-term tissue culture. *Cancer Res* 36: 4595-4601, 1976.
21. Dupont J, Karas M and LeRoith D: The potentiation of estrogen  
 on insulin-like growth factor I action in MCF-7 human breast  
 cancer cells includes cell cycle components. *J Biol Chem* 275:  
 35893-35901, 2000.
22. Zhu J, Li X, Kong X, *et al*: Testin is a tumor suppressor and  
 prognostic marker in breast cancer. *Cancer Sci* 103: 2092-2101,  
 2012.
23. Altucci L, Addeo R, Cicatiello L, *et al*: 17beta-estradiol induces  
 cyclin D1 gene transcription, p36D1-p34cdk4 complex activa-  
 tion and p105Rb phosphorylation during mitogenic stimulation  
 of G (1)-arrested human breast cancer cells. *Oncogene* 12: 2315,  
 1996.
24. Kodama F, Greene GL and Salmon SE: Relation of estrogen  
 receptor expression to clonal growth and antiestrogen effects on  
 human breast cancer cells. *Cancer Res* 45: 2720-2724, 1985.
25. Lu J, Pierron A and Ravid K: An adenosine analogue, IB-MECA,  
 down-regulates estrogen receptor  $\alpha$  and suppresses human  
 breast cancer cell proliferation. *Cancer Res* 63: 6413-6423,  
 2003.
26. Wormke M, Stoner M, Saville B, *et al*: The aryl hydrocarbon  
 receptor mediates degradation of estrogen receptor  $\alpha$  through  
 activation of proteasomes. *Mol Cell Biol* 23: 1843-1855, 2003.
27. Wijayaratne AL and McDonnell DP: The human estrogen  
 receptor- $\alpha$  is a ubiquitinated protein whose stability is affected  
 differentially by agonists, antagonists, and selective estrogen  
 receptor modulators. *J Biol Chem* 276: 35684-35692, 2001.
28. Horwitz KB, Koseki Y and McGuire WL: Estrogen control of  
 progesterone receptor in human breast cancer: role of estradiol  
 and antiestrogen. *Endocrinology* 103: 1742-1751, 1978.
29. Dauvois S, White R and Parker MG: The antiestrogen ICI  
 162675 disrupts estrogen receptor nucleocytoplasmic shuttling.  
*J Cell Sci* 106: 1377-1388, 1993.
30. Baker P, Wilton J, Jones C, Stenzel D, Watson N and Smith G:  
 Bile acids influence the growth, oestrogen receptor and  
 oestrogen-regulated proteins of MCF-7 human breast cancer  
 cells. *Br J Cancer* 65: 566, 1992.
31. Horwitz KB, Mockus MB and Lessey BA: Variant T47D human  
 breast cancer cells with high progesterone-receptor levels  
 despite estrogen and antiestrogen resistance. *Cell* 28: 633-642,  
 1982.
32. Rubio M, Werbajh S, Cafferata E, *et al*: TNF-alpha enhances  
 estrogen-induced cell proliferation of estrogen-dependent  
 breast tumor cells through a complex containing nuclear factor-  
 kappa B. *Oncogene* 25: 1367-1377, 2005.

- 1 33. Fisher B, Costantino JP, Wickerham DL, *et al*: Tamoxifen for  
2 prevention of breast cancer: report of the National Surgical  
3 Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer*  
4 *Inst* 90: 1371-1388, 1998. 61
- 5 34. Ali S and Coombes RC: Endocrine-responsive breast cancer  
6 and strategies for combating resistance. *Nat Rev Cancer* 2:  
7 101-112, 2002. 62
- 8 35. Yao K and Jordan VC: Questions about tamoxifen and the  
9 future use of antiestrogens. *Oncologist* 3: 104-110, 1998. 63
- 10 36. Cortesi L, De Matteis E, Rashid I, *et al*: Distribution of second  
11 primary malignancies suggests a bidirectional effect between  
12 breast and endometrial cancer: a population-based study. *Int J*  
13 *Gynecol Cancer* 19: 1358, 2009. 64
- 14 37. Gutierrez MC, Detre S, Johnston S, *et al*: Molecular changes  
15 in tamoxifen-resistant breast cancer: relationship between  
16 estrogen receptor, HER-2, and p38 mitogen-activated protein  
17 kinase. *J Clin Oncol* 23: 2469-2476, 2005. 65
- 18 38. Levin ER: Bidirectional signaling between the estrogen receptor  
19 and the epidermal growth factor receptor. *Mol Endocrinol* 17:  
20 309-317, 2003. 66
- 21 39. Osborne C, Wakeling A and Nicholson R: Fulvestrant: an  
22 oestrogen receptor antagonist with a novel mechanism of  
23 action. *Br J Cancer* 90: S2-S6, 2004. 67
- 24 40. Buijs C: Long-term side effects of adjuvant breast cancer  
25 treatment. University Library of Groningen[Host], 2008. 68
- 26 41. Klinge CM: Estrogen receptor interaction with co-activators  
27 and co-repressors. *Steroids* 65: 227-251, 2000. 69
- 28 42. Hsu SM, Chen YC and Jiang MC: 17 $\beta$ -estradiol inhibits tumor  
29 necrosis factor- $\alpha$ -induced nuclear factor- $\kappa$ B activation by  
30 increasing nuclear factor- $\kappa$ B p105 level in MCF-7 breast cancer  
31 cells. *Biochem Biophys Res Commun* 279: 47-52, 2000. 70
- 32 71
- 33 72
- 34 73
- 35 74
- 36 75
- 37 76
- 38 77
- 39 78
- 40 79
- 41 80
- 42 81
- 43 82
- 44 83
- 45 84
- 46 85
- 47 86
- 48 87
- 49 88
- 50 89
- 51 90
- 52 91
- 53 92
- 54 93
- 55 94
- 56 95
- 57 96
- 58 97
- 59 98
- 60 99
- 100
- 101
- 102
- 103
- 104
- 105
- 106
- 107
- 108
- 109
- 110
- 111
- 112
- 113
- 114
- 115
- 116
- 117
- 118
- 119
- 120